



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

of oxalic acid upon naphthylamine with that procured by treating a naphthalin-sulphate with cyanide of potassium. M. V. Merz had found the fusing-point of the latter acid to be 140° , whilst for the former I had observed the fusing-point 160° . M. O. Olshausen has since prepared in my laboratory a quantity of cyanide of naphthyl according to Merz's process. The acid obtained from this cyanide by treatment with an alkali, thrice recrystallized and finally purified by distillation, was likewise found to fuse exactly at 160° . Menaphtoxylamide, procured from the same source, exhibited the fusing-point 203° , while the compound I had formerly examined fused at 204° *. The identity of the acids obtained by the two processes is thus satisfactorily established.

- VI. "Account of some recent Observations on Sun-spots, made at the Kew Observatory." By WARREN DE LA RUE, Esq., F.R.S., BALFOUR STEWART, Esq., F.R.S., and BENJAMIN LOEWY, Esq., F.R.A.S. Received June 2, 1868.

(Abstract.)

The authors, after reviewing briefly the two theories on the nature of sun-spots, which are still subjects of dispute, refer to the stereoscopic views obtained and the results published in their 'Researches on Solar Physics,' and state the reasons which have led them to believe that sun-spots are cavities and at a lower level than the sun's photosphere. Their opinion has been recently strengthened by observations of a sun-spot on the 7th of May, which in disappearing produced in two successive photograms indentations in the west limb.

After proving by the measurements made, which, with the calculations, are appended to their paper, that there can be no doubt about the identity of the heliographical elements of the previously observed spot and the successive indentations, they prove from the established details of the phenomena of sun-spots that such indentations must under all circumstances be very rare occurrences, and state fully the conditions favourable to the recurrence of similar observations, inviting observers to give their particular attention to them.

- VII. "The Formation and Early Growth of the Bones of the Human Face." By GEORGE W. CALLENDER, Lecturer on Anatomy at St. Bartholomew's Hospital. Communicated by JAMES PAGET, F.R.S. Received June 2, 1868.

(Abstract.)

These notes refer to some few points with which we are as yet imper-

* The fusing-point of this substance is, by misprint (Proc. Roy. Soc. vol. xvi. p. 302), stated to be 244° , instead of 204° .—A. W. H.